

RÉFÉRENCES BIBLIOGRAPHIQUES

Abecia L., Mcewan N. R., Newbold C. J., Fondevila M., Balcells J. (2004). Molecular profiling of the major bacterial species in the rabbit caecum as affected by therapeutical doses of antibiotics. In: Proceedings of the 8th World Rabbit Congress. *World Rabbit Science Association*. Paris: 404-409

Akolkar A. V. (2009). Isolation and Characterization of Halophilic Archaea: Production, Characterization and Application of Extracellular Protease from *Halobacterium* sp. SP1(1).Thèse de Doctorat en microbiologie .Université de Baroda. Inde.

Amoozegar M. A., Malekzadeh F., Malik K.A. (2003). Production of amylase by newly isolated moderate halophile, *Halobacillus* sp. strain MA-2. *Microbiol Methods* 52: 353-359.

Amziane M., Djenane Z., Yaici L., Remane S., Selama O., Zerrouki A., Madour N., Sadouk Z., Bechet M., Chollet M., Jacques P., Abderrahmani A., Nateche F. (2012). Mise en évidence du potentiel en molécules bio-actives à partir de souches de *Bacillus* halotolérantes isolées des lacs salés du sud algérien. *J. hydrocarb. Mine environ. Res.* 3(2).

Ayad R. (2011). Screening d'activités hydrolytiques extracellulaires chez des microorganismes halophiles aérobies isolés d'environnements hypersalins de l'Est algérien. Mémoire de Magister en Sciences Alimentaires Option: Biotechnologie Alimentaire. Institut de la Nutrition, de l'Alimentation et des Technologies Agro-Alimentaires (I.N.A.T.A.A).Université de Constantine. Algérie.

Azandegbe A. (2010). Etude de la structure des communautés bactériennes du sédiment et de l'écologie de *Vibrio aestuarianus* pathogène de l'huître creuse *Crassostrea gigas* dans deux sites ostréicoles. Thèse de Doctorat en Microbiologie. Université de Brest, France.

Baghdad Belhadj-Semmar F. Z. (2012). Etude du stress osmotique, acide et thermique chez des Lactobacilles. Mémoire de Magister en Biotechnologie Option Génie microbiologique. Université d'Oran, Algérie.

Barkat S., Hoffmann L., Boumezbeur A. (2004). ATLAS [IV] des zones humides Algériennes d'importance internationale. Ed : Direction Générale des Forêts. Ministère de l'Agriculture et du Développement Rural.

Bechtel incorporated (USA) (1975). Développement des ressources en eau et de l'Agriculture. Région de Sétif. Etude préliminaire et conditions de réalisation – volume II. In : Barkat S., Hoffmann L., Boumezbeur A. (2004). ATLAS [IV] des zones humides Algériennes d'importance internationale. Ed : Direction Générale des Forêts. Ministère de l'Agriculture et du Développement Rural.

Berrada I., Willems A., De Vos P., El fahime E., Swings J., Bendaou N., Melloul M., Amar M. (2012). Diversity of culturable moderately halophilic and halotolerant bacteria in a marsh and two salterns a protected ecosystem of Lower Loukkos (Morocco). *African Journal of Microbiology Research* 6(10): 2419-2434.

Berrada I. (2012). Etude telluriques e la biodiversité des bactéries tellurique halophiles au niveau du BAS LOUKKOS (Marais Salants et Marecage de BEGGARA).Thèse de Doctorat en microbiologie et biologie moléculaire. Université Mohammed V –Agdal, Faculté Des Sciences. Rabat.

Bonneté F., Ebel C., Eisenberg H., Zaccai G. (1993). Biophysical study of halophilic malate deshydrogenase in solution: revised subunit structure and solvent interactions in native and recombinant enzyme. *J. Chem. soc. Faraday Trans* 89(15): 2659-66.

Boutaiba S, Bhatnagar T, Hacene H, Mitchell DA, Baratti JC (2006). Preliminary characterization of a lipolytic activity from an extremely halophilic archaeon, *Natronococcus sp.* *J. Mol. Catal. B Enzym.* 41: 21-26.

Boutaiba S., Hacene H., Bidle K.A., Maupin-Furlow J.A. (2011). Microbial diversity of the hypersaline Sidi Ameur and Himalatt Salt Lakes of the Algerian Sahara. *J Arid Environ* 75: 909-916.

Bowman J. P. (2007). Bioactive Compound Synthetic Capacity and Ecological Significance of Marine Bacterial Genus *Pseudoalteromonas*. *Mar Drugs* 5(4): 220-241.

Cánovas D., Vargas C., Kneip S., Morón M. J., Ventosa A., Bremer E., Nieto J. J. (2000). Genes for the synthesis of the osmoprotectant glycine betaine from choline in the moderately halophilic bacterium *Halomonas elongata* DSM 3043, USA. *PubMED*:455-63.

Caumette P., Cohen Y., Matheson R. (1991). Isolation and characterization of *Desulfovibrio halophilus* sp. Nov. A halophilic sulphate-reducing bacteria isolated from Solar Lake (Sinai). *Systematic and Applied Microbiology* 14 : 33-38.

Caumette P. (1998). Les bactéries halophiles: la vie dans les conditions extrêmes de salinité. *C R Acad Agric Fr.* 84: 11-21.

Cayol J. L., Ollivier B., Alazard D., Amils R., Godfroy A., Marty D., Piette F., Prieur D. (2012). Les conditions de vie extrêmes sur la planète et exobiologie. In : *Bertrand J.C.* (ed.), Caumette P. (ed.), Lebaron P. (ed.), Matheron R. (ed.), Normand P. (ed.) *Ecologie microbienne : microbiologie des milieux naturels et anthropisés.* Pau : PUPPA : 363-371.

Chedad K., Assobhei O. (2007). Etude de la survie des bactéries de contamination fécale (coliformes fécaux) dans les eaux de la zone ostréicole de la lagune de Oualidia (Maroc). *Bulletin de l Institut Scientifique, Rabat, section Sciences de la Vie* 29 : 71 -79.

Chena W., Changb J., Chiua C., Changc S. , Chenaand W. and Jianga C. (2005). *Caldimonas taiwanensis* sp. nov., a amylase producing bacterium isolated from a hot spring. *Syst. Appl. Microbiol.* 28 : 415–420.

Claus D., Berkeley R. C. W. (1986). Genus *Bacillus* Cohn 1872. In : Sneath PHA (ed) *Bergey's Manual of Systematic Bacteriology*, vol 2. Williams and Wilkins, Baltimore :1105-1139.

Cojoc R., Merciu S., Popescu G., Dumitru L., Kamekura M., Enache M. (2009). Extracellular hydrolytic enzymes of halophilic bacteria isolated from a subterranean rock salt crystal. *Rom. Biotechnol. Lett.* 14:4658- 4664.

Copin –Montégut G. (1996). Chimie de l'eau de mer. Institut océanographique, Paris. In : Boutaiba S., Hacene H., Bidle K.A., Maupin-Furlow J.A. (2011). Microbial diversity of the hypersaline Sidi Ameer and Himalatt Salt Lakes of the Algerian Sahara. *J Arid Environ* 75: 909-916.

Costenaro L. (2001). Interactions faibles protéine –protéine en solution: La malate déshydrogénase halophile. Thèse de Doctorat en Cristallographie et RMN biologique (physique). Université Joseph Fourier –Grenoble 1. France.

Dang H., Zhu H., Wang J., Li T. (2009). Extracellular hydrolytic enzyme screening of culturable heterotrophic bacteria from deep-sea sediments of the Southern Okinawa Trough. *World J. Microbiol. Biotechnol* 25: 71-79.

Dauga C., Dore J., Sghir A. (2005). La diversité Insoupçonnée du monde microbien. *MEDECINE/SCIENCES* 21 : 290-6.

Djaballah Ch. (2010). Biodiversité des Actinomycètes halophiles et halotolérants isolés de la Sebkhha de Ain M'LILA. Mémoire de Magister en Microbiologie .option : écologie microbienne. Université Mentouri Constantine ALGERIE.

Dohrmann A. B., Müller V. (1999). Chloride dependence of endospore germination in *Halobacillus halophilus*. *Arch Microbiol* 172: 264-267.

Dollhopf S. L., Hashsham S. A., Dazzo F. B., Hickey R. F., Criddle C. S., Tiedje J. M. (2001). The impact of fermentative organisms on carbon flow in methanogenic systems under constant low-substrate conditions. *Appl Microbiol Biotechnol* 56 : 531-538.

Dorigo U., Volatierb L., Humberta J.F. (2005). Molecular approaches to the assessment of biodiversity in aquatic microbial communities. *Journal of Water Research. Elsevier* : 0043-1354.

Drancourt M., Bollet C., Carlizoz A., Martelin R., Gayral J.-P., Raoult D. (2000). 16S ribosomal DNA sequence analysis of a large collection of environmental and clinical unidentifiable bacterial isolates. *J. Clin. Microbiol.* 38: 3623–3630.

Edgerton M. E., Brimblecome P. (1981). Thermodynamics of halobacterial environments. *Can J Microbiol* 27: 899-909.

Elevi R., Assa P., Birbir M., Ogan A., Oren A. (2004). Characterization of extremely halophilic archaea isolated from the Ayvalik salterns, Turkey. *World J. Microbiol. Biotechnol.* 20: 719- 725.

Faugier A. (2010). Diversité bactérienne des sols: accès aux populations à effectifs minoritaires “the rare biosphère”. Thèse de Doctorat. Ecole doctorale Electronique, Electrotechnique, Automatique de Lyon. France.

Fischer S.G., Lerman L. S. (1983). DNA fragments differing by single base-pair substitutions are separated in denaturing gradient gels: Correspondence with melting theory. *Proc. NatL Acad. Sci.* 80 : 1579-1583.

Fisher M. M., Triplett E. W. (1999). Automated approach for ribosomal intergenic spacer analysis of microbial diversity and its application to freshwater bacterial communities. *Appl Environ Microbiol* 65: 4630-4636.

Garabito M. J., Marquez M. C., Ventosa A. (1998). Halotolerant Bacillus diversity in hypersaline environment. *Can J Microbiol.* 44: 95-102.

Gavrieli I. (1997). Halite deposition in the Dead Sea :1960-1993. In: Boutaiba S., Hacene H., Bidle K.A., Maupin-Furlow J.A. (2011). Microbial diversity of the hypersaline Sidi Ameur and Himalatt Salt Lakes of the Algerian Sahara. *J Arid Environ* 75: 909-916.

Gerhardt P., Murray R. G. E., Wood W. A., Krieg N. R. (1994). Methods for General and Molecular Bacteriology. Washington, DC. *American Society for Microbiology* 791.

Ghasemi Y., Rasoul-Amini S., Ebrahiminezhad A., Kazemi A., Shahbazia M., Talebniaa N. (2011). Screening and Isolation of Extracellular Protease Producing Bacteria from the Maharloo Salt Lake. *Iran. J. Pharm. Sci.* 7:175-180.

Gonzalez C., Gutierrez C., Ramirez C. (1978). *Halobacterium vallismortis* sp. nov. an amyolytic and carbohydrate-metabolizing, extremely halophilic bacterium. *Can J Microbiol.* 24: 710–715.

Govender L., Naidoo L., Setati M.E. (2009). Isolation of hydrolase producing bacteria from Sua pan solar salterns and the production of endo-1,4-b-xylanase from a newly isolated haloalkaliphilic *Nesterenkonia* sp. *Afr. J. Biotechnol.* 8: 5458-5466. In : Mathabatha E. S. (2010). Diversity and industrial potential of hydrolaseproducing halophilic/halotolerant eubacteria. *African Journal of Biotechnology* 9 (11): 1555-1560.

Grant W. D., Kamekura M., McGenity T. J., Ventosa A. (2001). Class III. *Halobacteria* class nov. In: Boone D. R., Castenholz R.W., Garrity G.M (eds) Bergey’s manual of systematic bacteriology, vol 1, 2nd edn, The *Archaea* and the deeply branching and phototrophic *Bacteria*. *Springer* : 294–301.

Guesmi A., Ettoumi B., El Hidri D., Essanaa J., Cherif H. Mapelli F., Marasco R., Rolli E., Boudabouss A., Cherif A. (2013). Uneven Distribution of *Halobacillus trueperi* species in arid natural saline systems of southern Tunisian Sahara. *Environmental Microbiology (ed) Springer* 10.1007/s00248-013-0274-4.

Gupta R. Gupta N. Rathi. P. (2004). Bacterial lipases: an overview of production, purification and biochemical properties. *Appl Microbiol Biotechnol.* 64: 763–781

Hacène H., Rafa F., Chebhouni N., Boutaiba S., Bhatnagar T., Baratti J. C., Ollivier B. (2004). Biodiversity of prokaryotic microflora in El Golea salt Lake, Algerian Sahara. *J Arid Environ* 58: 273-284.

Hänelt I., Müller V. (2013). Molecular mechanism of adaptation of the moderately halophilic bacterium *Halobacillus halophilus* to its environment . *J of Life* 3: 234-243.

Hedi A., Sadfi N., Fardeau M-L., Rebib H., Cayol J-L., Ollivier B., Boudabous A. (2009). Studies on the Biodiversity of Halophilic Microorganisms Isolated from El-Djerid Salt Lake (Tunisia) under Aerobic Conditions. *Int J Microbiol:* 1-17.

Henze M., Van Loosdrecht M. C. M., Ekama G., Brdjanovic D. (2008). Biological wastewater treatment: principles, modelling and design. *Technol Eng:* 511.

Hrabak O. (1992). Industrial production of polyhydroxybutyrate. *FEMS Microbiology Rev.* 103: 251-256.

Huybens N., Mainil J., Marlier D. (2009). Les techniques de biologie moléculaire d'analyse des populations bactériennes complexes. *Annales de Médecine Vétérinaire* 153: 112-128.

Imhoff J. J., Sahl H., Soliman G., Truper H. (1979). The wadi Natrun: chemical composition and microbial mass developments in alkaline brines of eutrophic desert lakes. *Geomicrobiology journal* 1:219-234. In: Boutaiba S., Hacene H., Bidle K.A., Maupin-Furlow J.A. (2011). Microbial diversity of the hypersaline Sidi Ameur and Himalatt Salt Lakes of the Algerian Sahara. *J Arid Environ* 75: 909-916.

Joffin J.-N. et Leyral G. (2006). Microbiologie Technique .Tome 1.Dictionnaire des Techniques 4 éme édition. Collection biologie technique .les éditions Scérène. CRDP. Aquitaine.

Johnson A. M., Thurlow L. R., Zwenger R. S. and Gillock E. T. (2007). Partial Characterization of Two Moderately Halophilic Bacteria from a Kansas Salt Marsh. *The Prairie Naturalist* 39(1): 29–39.

Kamekura M., Hamakawa T., Onishi H. (1982). Application of halophilic nuclease H of *Micrococcus varians* *subsp. halophilus* to commercial production of flavoring agent 5'-GMP. *Appl Environ Microbiol* 44: 994–995.

Kamekura M., Dyall-Smith M. L. (1995). Taxonomy of the family *Halobacteriaceae* and the description of two new genera *Halorubrobacterium* and *Natrialba*. *J. Gen. Appl. Microbiol.* 41:333-350.

- Kamekura M. (1998).** Diversity of extremely halophilic bacteria. *Extremophiles* 2: 289-295.
- Kharaka Y. K., Hanor J. S. (2005).** 5.16 - Deep Fluids in the Continents: I. Sedimentary Basins . *Treatise on Geochemistry* 5(16): 499-540.
- Kharroub K. (2007).** Identification et étude moléculaire des bactéries et des archéobactéries aérobies halophiles de la sebkha Ezzemoul (Ain M'Lila). Thèse de doctorat, Université Mentouri-Constantine.
- Kaurichev, I. S. (1980).** Practicas de Edofologia. Mir Publishers, Moscow.
- Khunt M., Pandhi N., Rana A. (2011).** Amylase from moderate halophiles isolated from wild ass excreta. *Int. J. Pharm. Bio. Sci.* 1: 586- 592.
- Kitouni M. (2007).** Isolement de bactéries Actinomycétales productrices d'antibiotiques à partir d'écosystèmes extrêmes. Identification moléculaire des souches actives et caractérisation préliminaire des substances élaborées. Thèse de Doctorat d'État en Microbiologie appliquée. Université Mentouri Constantine. Algérie.
- Kloppmann W., Bourhane A., Asfirane F. (2011).** Méthodologie de diagnostic de l'origine de la salinité des masses d'eau. Emploi des outils géochimiques, isotopiques et géophysiques. BRGM :Giosciences pour une terre durable. BRGM/RP-60026-FR
- Klouche K. N., Aissaoui N., Nas F., Cayol J. L., Ghellai L. (2013).** A Novel Halotolerant Bacterium Isolated from El Goléa Lake in Algeria and Antimicrobial Potential of this Strain. *Journal of Agricultural Science and Technology*: 825-834.
- Lefebvre O. (2005).** Application des micro-organismes halophiles au traitement des effluents industriels hypersalins. Thèse de doctorat en sciences et procédés biologiques et industriels, Spécialité Génie des procédés, Ecole Nationale Supérieure Aronomique, Montpellier. France.
- Litchfield C. D., Gillevet P. M. (2002).** Microbial diversity and complexity in Hypersaline environments: A preliminary assessment. *J. Ind. Microbiol & Biotechnol* 28 (1): 48-55.
- Logue J. B., BÜrgmann H., Robinson C.T. (2008).** Progress in the ecological genetics and biodiversity of freshwater bacteria. *Bioscience* 58(2):103-113.
- Longeon A., Peduzzi J., Barthelemy M., Corre S., Nicolas J. L., Guyot M. (2004).** Purification and partial identification of novel antimicrobial protein from marine bacterium *Pseudoalteromonas* species strain X153. *Mar Biotechnol* 6: 633-641. *stère de l'Agriculture et du Développement Générale des Forêts.*
- Ma Y., Galinski E. A., Grant W. D., Oren A., Ventosa A. (2010).** Life in Saline Environments. *Applied and Environmental Microbiology*: 6971–6981.

McHardy A. C., Rigoutsos I. (2007). What's in the mix: phylogenetic classification of metagenome sequence samples. *Curr Opin Microbiol.* 10: 499-503.

Madern D., Pfisler C., Zaccai G. (1995). Mutation at a single acidic aminoacid enhances the halophilic behaviour of malate deshydrogenase from *Haloarcula marisimortui* in physiological salts. *Eur. J. Biochem* 230(3):1088-95.

Madigan M.T., Martinko J. M. (2007). Brock Biology of Microorganisms, 11th edition. Pearson Prentice Hall, Upper Saddle River, New Jersey.

Mariana N. G., Virginia A. V., Henry A., Michal S. (2008). Lipolytic enzyme production by halophilic/halotolerant microorganisms isolated from laguna verde, Bolivia, *revista boliviana de quimica* 25:1.

Marzorati M., Alma A., Sacchi L., Pajoro M., Palermo S., Brussetti L., Raddadi N., Balloi A., Tedeschi R., Clementi E., Corona S., Quaglino F., Bianco P., Beninati T., Bandi C., Daffonchio D. (2006). A Novel *Bacteroidetes* Symbiont Is Localised in *Scaphoideus titanus*, the Insect Vector of Flavescence Dorée in *Vitis vinifera*. *Appl Environ Microbiol* 72(2):1467-1475.

Montalvo-Rodríguez R., Vreeland R. H., Oren A., Kessel M., Betancourt C., López-Garriga J. (1998). *Halogeometricum borinquense* gen. nov., sp. nov., a novel halophilic archaeon from Puerto Rico. *Int. J. Syst. Bacteriol.* 48: 1305–1312.

Moreno M. L., García M. T., Ventosa A., Mellado E. (2009). Characterization of *Salicola* sp. IC10, lipase- and protease-producing extreme halophile. *FEMS Microbiol. Ecol.* 68: 59-71.

Moreno M. L., Piubeli F., Bonfá M. R., García M. T., Durrant L. R., Mellado E. (2013). Analysis and characterization of cultivable extremophilic hydrolytic bacterial community in heavy-metal-contaminated soils from the Atacama Desert and their biotechnological potentials. *J. Appl. Microbiol.* 113: 550- 559.

Müller-Santos M., de Souza E. M., Pedrosa F. D., Mitchell D. A., Longhi S., Carriere F., Canaan S., Krieger N. (2009). *Biochim Biophys Acta.* 1791(8): 719–729.

Muyzer G., Dewaal E. C., Uitterlinden A. G. (1993). Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction - amplified genes coding for 16S ribosomal RNA. *Appl Environ Microbiol* 59: 695-700.

Muyzer G., Smalla K. (1998). Application of denaturing gradient gel electrophoresis (DGGE) and temperature gradient gel electrophoresis (TGGE) in microbial ecology. *Antonie Van Leeuwenhoek International Journal of General and Molecular Microbiology* 73: 127-141.

Oh B. C., Kim H. K., Lee J. K., Kang S. C., Oh T. K. (1999). *Staphylococcus haemolyticus* lipase: biochemical properties, substrate specificity and gene cloning. *FEMS Microbiol Lett* 179: 385–392.

Ollivier B., Caumette P., Garcia J. L., Mah R. (1994). Anaerobic bacteria from hypersaline environments. *Microbiol Rev* 58: 27-38.

Oren A. (1994). The ecology of the extremely halophilic archaea. *FEMS Microbiol rev* 13: 415-440.

Oren A. (1999). The enigma of square and triangular bacteria. In: Seckbach J(Ed.), *Enigmatic Microorganisms and life in Extreme Environmental Habitats. Kluwer Academic Publishers, Dordrecht: 337-355.*

Oren A. (2000). Salts and brines. In: Whitton BA and M Potts (Eds.), *Ecology of Cyanobacteria: Their Diversity in Time and Space. Kluwer Academic Publishers, Dordrecht: 281– 306.*

Oren A., Rodriguez-Valera F. (2001). The contribution of *Salinibacter ruber* species to the red coloration of saltern crystallizer ponds. *FEMS Microbiol Ecol* 36: 123-130.

Oren A. (2002). Diversity of halophilic microorganisms: environments phylogeny, physiology, and applications. *J Ind Microbiol Biotechnol* 28: 56-63.

Oren A. (2006). Life at high salt concentrations. In: Dworkin M., Falkow S., Rosenberg E., Schleifer K. H., Stackebrandt E (eds) *The Prokaryotes. A Handbook on the Biology of Bacteria: Ecophysiology and Biochemistry. Springer 2: 263-282.*

Oren A. (2008). Microbial life at high salt concentrations: phylogenetic and metabolic diversity. *Saline Syst* 4(2): 1746-1448.

Oren A. (2010). Industrial and environmental applications of halophilic microorganisms. *Environmental Technology* 31: 8–9, 825–834.

Oren A., Ventosa A. (2013). Subcommittee on the taxonomy of *Halobacteriaceae* and subcommittee on the taxonomy of *Halomonadaceae*. *Int. J. of Syst and Evol Microbiology* 63: 3540–3544.

Paul A., Hackbarth S., Vogt R. D., Roeder B., Burnison B. K., Steinberg C. E. W. (2004). Photogeneration of singlet oxygen by humic substances: comparison of humic substances of aquatic and terrestrial origin. *Photochem Photobiol Sci* 3: 273–280.

Pikuta E. V., Hoover R. B. (2007). Microbial Extremophiles at the Limits of Life. *Critical Reviews in Microbiology* 33:183–209.

Pinaki S., Ekramul I. (2012). Metagenomic Approaches in Microbial Bioremediation of Metals and Radionuclides. In: T. Satyanarayana et al. (eds.), *Microorganisms in Environmental Management: Microbes and Environment* 978: 94-007.

Post F. (1981). Microbiology of the Great Salt Lake north arm. *Hydrobiologia* 81: 59-69.

Pastor J. M., Manuel S. , Argandoña M., Bernal V., Reina-Bueno M., Csonka L. N., Iborra J. L., Vargas C., Nieto J. J., Cánovas M. (2010). Ectoines in cell stress protection: Uses and biotechnological production. *Elsevier:782-801.*

Richard S. B., Madern D., Garcin E., Zaccai G. (2000). Halophilic adaptation: novel solvent protein interactions observed in the 2.9 and 2.6 Å resolution structures of the wild type and a mutant of malate dehydrogenase from *Haloarcula marismortui*. *Biochemistry* 39: 992-1000

Riesenfeld C. S., Schloss P. D., Handelsman J. (2004). Metagenomics: genomic analysis of microbial communities. *Annu. Rev. Genet.* 38: 525–552. In: T. Satyanarayana et al. (2012). *Microorganisms in Environmental Management: Microbes and Environment*. Springer: 978-94-007.

Rodier J., Legube B., Merlet N. et coll. (2009). L'analyse de l'eau. Les éditions Dunod, 9ème édition.

Rodriguez-Valera F. (1993). Introduction to saline environments. In: Vreeland R. H., Hochstein L.I (eds) The biology of halophilic bacteria. *CRC Press Boca Ratom FL*: 1-23.

Roessler M., Müller V. (1998). Quantitative and physiological analyses of chloride dependence of growth of *Halobacillus halophilus*. *Appl. Environ. Microbiol* 64: 3813-3817.

Roessler M., Müller .V. (2002). Chloride, a new environmental Signal molecule involved in gene regulation in a moderately halophilic bacterium, *Halobacillus halophilus*. *J. Bacteriol.*184: 6207-6215.

Rohban R., Amoozegar M. A., Ventosa A. (2009). Screening and isolation of halophilic bacteria producing extracellular hydrolyses from Howz Soltan Lake. *Iran. Ind. Microbiol. Biotechnol.* 36: 333-340. In : Mathabatha E. S. (2010). Diversity and industrial potential of hydrolaseproducing halophilic/halotolerant eubacteria. *African Journal of Biotechnology* 9 (11): 1555-1560.

Sánchez-Porro C., Martin S., Mellado E., Ventosa A. (2003). Diversity of moderately halophilic bacteria producing extracellular hydrolytic enzymes. *Appl. Microbiol.* 94: 295-300.

Sass A. M., McKew B. A., Sass H., Fichtel J., Timmis K. N., McGenity T. J. (2008). Diversity of Bacillus-like organisms isolated from deep-sea hypersaline anoxic sediments. *Saline Systems* 4 (8): 1-11. 10. 1186/1746-1448-4-8.

Saum S. H., Müller V. (2008). Régulation of osmoadaptation in the moderate halophile *Halobacillus halophilus* : Chloride,glutamate and Switching osmolyte strategies. *Saline Syst*: 1746-1448-4-4.

Shendure J., Ji H. (2008). Next-generation DNA sequencing. *Nat Biotechnol.* 26: 1135-1145.

Sivakumar U., Boutaiba S., Matthew A. H., Julie A. M.-F. (2009). LccA, an Archaeal Laccase Secreted as a Highly Stable Glycoprotein into the Extracellular Medium by *Haloferax volcanii*. *Applied and Environmental Microbiology* : 733–743.

Sleator R. D., Hill C. (2001). Bacterial osmoadaptation: the role of osmolytes in bacterial stress and virulence. *FEMS Microbiology Reviews* 26: 49-71.

Slimani R. (2006). Contribution à l'évaluation d'indicateurs de pollution environnementaux dans la région de Ouargla: Cas des eaux de rejets (agricoles et urbaines). Mémoire de Magister Spécialité : Agronomie Saharienne Option : Protection des écosystèmes en zones arides. Université Kasdi Merbah. Ouargla.

Stackebrandt E., Goebel B. M. (1994). Taxonomic note: A place for DNA-DNA reassociation and 16SrDNA sequence analysis in the present species definition in bacteriology. *Int. jour. sys. Bacteriol* 44:846-849. Streptomyces to sodium chloride as a taxonomic tool. *Appl. Microbiol.* 16: 1134- 1136.

Torsvik V., Øvreås L. (2008). Microbial Diversity, Life Strategies, and Adaptation to Life in Extreme Soils. in: P. Dion and C.S. Nautiyal (eds.). Microbiology of Extreme Soils. *Soil Biology 13. Springer:* 15-43.

Tsai Y.-L., Olson B. H. (1991). Rapid method for direct extraction of DNA from soil and sediments. *App Environ Microbiol* 57: 1070-1074.

Ventosa A., Mellado E., Sanchez-Porro C., Marquez M. C. (2008). Halophilic and halotolerant Micro-Organisms from Soils. In : P. Dion and C.S. Nautiyal (eds.). Microbiology of Extreme Soils. *Soil Biology 13. Springer:* 87–115.

Whitman William B., Parte A., Goodfellow M., Kämpfer P., Busse H.-J., Trujillo Martha E., Ludwig W., Suzuki K.-I. (2012). Bergey's Manual of Systematic Bacteriology: The Actinobacteria, Volume 5, Parties 1 à 2. *Springer:* 1750.

Woese C. R. (1987). Bacterial evolution. *Microbial Rev* 51, 221-271.

Won-Jae Ch., Park Da Y., Jeong S.-Ch., Chang Y.-K., Kwang H. S. (2011). Isolation and Characterization of Starch-hydrolyzing Pseudoalteromonas sp. A-3 from the Coastal Sea Water of Daecheon, Republic of Korea. *Korean. J. Microbiol. Biotechnol.* 39 : 4, 317–323.

Woodward J., Wiseman A. (1984). Topics in enzyme and fermentation biotechnology, Volume 8. New York: *John Wiley & Sons:* 9–30.

Yachai M. (2009). Carotenoid production by halophilic Archaea and its applications. Thesis of Doctorat, university Prince of Songkla. 173.

Yanan W., Zhiying P., Shuangjiang L. (2004). Ecological study of Bacillus in sediments of marine-cultural pond. *Journal of Zhanjiang Ocean University* 24(4): 23-27.

Zhilina T. N., Zavarzin G. A. (1990). Anew extremely halophilic homoacetogen bacteria *Acetohalobium arabaticum*, gen. nov. sp. nov. *Dokl. Akad. Nauk SSSR.* 311: 745-747.